

MYC-Tag Mouse mAb

www.dia-an.cn


Cat#: 2097

027-59325199

sevice@dia-an.com

Only For Research. Not For Diagnosis.

Synonyms:	EQKLISEEDL
Attribute:	Mouse monoclonal Antibody
Isotype:	IgG
Purity:	Antigen Affinity Purification
Application:	ELISA, WB, IP, IF
Calculated MW:	N/A
Observed MW:	N/A
Reactivity:	Human, Mouse, Rat, Zebra fish
Immunogen:	Peptide EQKLISEEDL
Buffer:	PBS with 0.1% sodium azide and 50% glycerol, pH 7.2
Storage:	Store at -20°C. Do not aliquot
Recommended Dilution:	WB: 1:2000-5000 IP: 3ug/sample IF: 1:100-200



Western Blotting of MYC-Tag fused recombinant protein with Anti-MYC Mouse monoclonal antibody at dilution of 1:5000.

Background:

The MYC tag, derived from the c-MYC protein, is a popular epitope tag for detecting the expression of recombinant proteins in yeast, bacteria, insect, and mammalian cell systems. It can be used for affinity chromatography, then used to separate recombinant, overexpressed protein from wild type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits.

We focus on precise protein quantification