

## HA-Tag Rabbit mAb

www.dia-an.cn

Cat#: 3063R

027-59325199


sevice@dia-an.com

Only For Research. Not For Diagnosis.

<b>Synonyms:</b>	YPYDVPDYA
<b>Attribute:</b>	Rabbit Monoclonal Antibody
<b>Isotype:</b>	IgG
<b>Purity:</b>	Antigen Affinity Purification
<b>Application:</b>	ELISA, WB, IP, IF
<b>Reactivity:</b>	Protein fused with HA tag
<b>Immunogen:</b>	Peptide YPYDVPDYA
<b>Buffer:</b>	PBS with 0.1% sodium azide and 50% glycerol, pH 7.2
<b>Storage:</b>	Store at -20°C. Do not aliquot
<b>Recommended Dilution:</b>	WB: 1:2000-1:10,000 IP: 3ug/sample IF: 1:100-1:200

### Background:

The HA tag is derived from the HA-molecule corresponding to amino acids 98-106. It has been extensively used as a general epitope tag in expression vectors. Many recombinant proteins have been engineered to express the HA tag, which does not appear to interfere with the bioactivity or the biodistribution of the recombinant protein.



Western Blotting of HA-Tag fused recombinant protein with Anti-HA Rabbit Monoclonal Antibody at dilution of 1:5000

We focus on precise protein quantification

    
Western blot of Zebra fish whole lysates  
with anti-H2A Rabbit Polyclonal Antibody  
at dilution of 1:1000